

SUPERSTABILITY AND FINITE TIME EXTINCTION FOR C_0 -SEMIGROUPS

D. CREUTZ, M. MAZO, JR., AND C. PREDA

ABSTRACT. A new approach to superstability and finite time extinction of strongly continuous semigroups is presented, unifying known results and providing new criteria for these conditions to hold analogous to the well-known Pazy condition for stability. That finite time extinction implies superstability which is in turn equivalent to several (both known and new) conditions follow from this new approach in a consistent fashion. Examples that the converse statements fail are constructed, in particular, an answer to a question of Balakrishnan on superstable systems not exhibiting finite time extinction.

1. INTRODUCTION

The study of C_0 -semigroups as a means to understand systems, particularly systems modeled by (partial) differential equations, has a long and rich history. A central concept in this study is the notion of exponential stability (often referred to simply as stability). A related stronger condition known as superstability has been the focus of much research (e.g. [Bal05], [Bal81], [NR92], [RW95], [Lum01]), in particular its connection with finite time extinction.

Balakrishnan [Bal05] poses the following: Are there physical (i.e. differential operator) examples of superstability which are not of type extinction-in-finite-time? We provide a positive answer to this question with a constructive example (in fact many similar examples can easily be obtained). We also collect, clarify and extend the existing results in the field with a new unified approach.

The notion of superstability first appeared (in a very rough form) in the seminal work of Hille and Philips [HP57] who were concerned primarily with the mathematical aspects of it and related properties, particularly the relationship between the spectrum of the infinitesimal generator and the stability of the semigroup. Later work refined and extended this notion, applying more complicated machinery ([NR92], [RW95]). More recently Balakrishnan [Bal05] (and others) have become interested in the superstability phenomena arising in the control theory of (physical) systems.

The key ingredient in our work is a new approach to the concepts of stability and superstability focusing on the “entry times” of the system into balls about the origin (in the Banach space). This approach, which has a certain probabilistic flavor though is not in itself probabilistic, allows us to unify the existing results in the field with (largely) new proofs. More importantly, we obtain analogues of certain well-known results about stability for superstable and finite-time-extinction systems. In particular, an analogue of Pazy’s condition [Paz83] for stability is given for both superstability and finite time extinction.

2. MAIN RESULTS

Our main result characterizing the types of stability is:

Theorem. *Let $\{T(t)\}$ be a C_0 -semigroup of bounded linear operators on a Banach space X . Define the relative entry time for each $r \in \mathbb{N}$ as*

$$u_r = \sup\{t_{r+1}(x) - t_r(x) : \|x\| \leq 1\} \quad \text{where} \quad t_r(x) = \inf\{t \geq 0 : \|T(t')x\| \leq e^{-r} \text{ for all } t' \geq t\}$$

Then

(i)	$\{T(t)\}$ is stable	if and only if	$\limsup_r u_r < \infty$
(ii)	$\{T(t)\}$ is superstable	if and only if	$\lim u_r = 0$
(iii)	$\{T(t)\}$ has finite time extinction	if and only if	$\sum_r u_r < \infty$

and our Pazy-type condition is:

Theorem. Let $\{T(t)\}$ be a C_0 -semigroup. Then, if for some $a > 0$,

- (i) $\int_a^\infty \|T(t)\|^p dt < \infty$ for some $0 < p < \infty$ then $\{T(t)\}$ is stable
- (ii) $\int_a^\infty |\log \|T(t)\||^{-p} dt < \infty$ for some $1 < p < \infty$ then $\{T(t)\}$ is stable
- (iii) $\int_a^\infty |\log \|T(t)\||^{-1} dt < \infty$ then $\{T(t)\}$ is superstable
- (iv) $\lim_{p \downarrow 0} \int_a^\infty |\log \|T(t)\||^{-p} dt < \infty$ then $\{T(t)\}$ has finite time extinction

3. PRELIMINARIES

A family $\{T(t)\}_{t \geq 0}$ of bounded linear operators on a Banach space X is called a **strongly continuous semigroup** (or **C_0 -semigroup**) when $T(0) = I$, $T(t+s) = T(t)T(s)$ for all $t, s \geq 0$, and $\lim_{t \downarrow 0} T(t) = I$ in the strong operator topology ($T(t)x \rightarrow x$ as $t \downarrow 0$ for all $x \in X$). As is well-known, this implies that the map $t \mapsto T(t)$ is (strongly) continuous.

For a strongly continuous semigroup $\{T(t)\}_{t \geq 0}$, define $\mathcal{D}$ to be the set of all $x \in X$ such that $\lim_{t \downarrow 0} t^{-1}(T(t)x - x)$ exists. The **infinitesimal generator** of the semigroup $\{T(t)\}_{t \geq 0}$ is the operator A on X , with the domain $D(A) = \mathcal{D}$, given by

$$Ax = \lim_{t \downarrow 0} \frac{T(t)x - x}{t}, \quad x \in D(A).$$

The name “infinitesimal generator” is justified by the fact that

$$Ax = \left. \frac{d(T(t)x)}{dt} \right|_{t=0}, \quad x \in D(A).$$

The pair $(A, \mathcal{D})$ and the semigroup $\{T_t\}_{t \geq 0}$ uniquely determine one another.

A strongly continuous semigroup $\{T(t)\}_{t \geq 0}$ is called **exponentially stable** (or just **stable**) when there exists constants $M > 0$ and $\rho > 0$ such that

$$\|T(t)\| \leq Me^{-\rho t} \quad \text{for all } t \geq 0.$$

Equivalently, define the **stability index** to be

$$\sup\{\nu > 0 : (\exists M > 0) \|T(t)\| \leq Me^{-\nu t} \text{ for all } t \geq 0\}$$

and stability is then the requirement that the index be positive.

The **growth characteristic** is

$$\omega_0 = \lim_{t \downarrow 0} \frac{\log \|T(t)\|}{t}$$

which, as is well-known, is equal to $-\nu$ where ν is the stability index (when the semigroup is stable).

It is then natural to define **superstability** to be the condition that the growth characteristic is $\omega_0 = -\infty$. Alternatively, superstability can be defined as the equivalent condition that the operators $T(t)$ be quasinilpotent (recall that an operator T is quasinilpotent when $\text{spec}(T) = \{0\}$).

A system is said to have **finite time extinction** when there is some $t_0 \geq 0$ such that $T(t)x = 0$ for all $t \geq t_0$ and all $x \in X$ with $\|x\| \leq 1$.

A semigroup $\{T(t)\}$ is **nilpotent** when there exists t_0 such that $T(t_0) = 0$. The smallest possible choice t_0 such that $T(t') = 0$ for all $t' > t_0$ is called the **index of nilpotency** for the semigroup.

The reader should note that in what follows we often defer the proofs until after all results are stated, the purpose being to stress the similarities among the theorems characterizing these three concepts (perhaps the most useful aspect of our approach).

4. FINAL ENTRY TIMES

Definition 1. Let $\{T(t)\}$ be a C_0 -semigroup on a Banach space X . For each $x \in X$ and $r \in \mathbb{N}$ the **final entry time** of x into the e^{-r} -ball is

$$t_r(x) := \inf\{t \geq 0 : \|T(t')x\| \leq e^{-r} \text{ for all } t' \geq t\}$$

and the **final entry time** of the 1-ball into the e^{-r} -ball (referred to from here on as just the final entry time of the e^{-r} -ball) is

$$t_r := \sup\{t_r(x) : \|x\| \leq 1\}$$

where we adopt the (usual) convention that the infimum of the empty set is ∞ (that is, $t_r(x) = \infty$ when no such time exists).

The **relative entry time** of x into the e^{-r} -ball is

$$u_r(x) := \begin{cases} t_{r+1}(x) - t_r(x) & \text{when } t_{r+1}(x) < \infty \\ \infty & \text{when } t_{r+1}(x) = \infty \end{cases}$$

and the **relative entry time** of the 1-ball into the e^{-r} -ball (referred to from here on as just the final entry time of the e^{-r} -ball) is

$$u_r := \sup\{u_r(x) : \|x\| \leq 1\}$$

Lemma 2. For any r we have $u_r = t_{r+1} - t_r$ (meaning when either or both $t_r, t_{r+1} = \infty$ then $u_r = \infty$).

Proof. First note that $u_r = \infty$ if and only if $t_{r+1} = \infty$ and that $t_r = \infty$ implies $t_{r+1} = \infty$ so we need only handle the case that all three are finite. By definition,

$$u_r + t_r = \sup\{u_r(x) : \|x\| \leq 1\} + \sup\{t_r(x) : \|x\| \leq 1\} \geq t_{r+1}.$$

On the other hand, there exists a sequence x_n such that $t_r(x_n) \rightarrow t_r$. So for any $\epsilon > 0$ and sufficiently large n we have $t_r(x_n) > t_r - \epsilon$ and so $t_r < t_r(x_n) + \epsilon$. By definition $t_{r+1} \geq t_{r+1}(x)$ for all x , in particular for the x_n . Then

$$t_{r+1} - t_r \geq t_{r+1}(x_n) - t_r(x_n) - \epsilon \geq u_r - \epsilon.$$

As ϵ is arbitrary, the claim follows. $\square$

Lemma 3. For any x and r with $\|x\| > e^{-r}$ we have $\|T(t_r(x))x\| = e^{-r}$. Moreover, $\|T(t_r)\| = e^{-r}$.

Proof. This follows directly from the strong continuity of $T(t)$ (which is automatic for right continuity and follows from the uniform boundedness principle for left continuity). $\square$

Lemma 4. The sequence u_r is nonincreasing in r : $u_{r+1} \leq u_r$. Hence the limit $\lim_r u_r$ always exists.

Proof. First note that if u_r ever attains ∞ then in fact $t_{r+1} = \infty$ at that point and for all subsequent t_r meaning that u_r remains ∞ from then on. So we need only handle the case when $u_r < \infty$ for all r (and therefore assume that $t_r < \infty$ for all r).

Now suppose that $u_r > u_{r-1}$ for some r . Then there is some x such that $u_r(x) > u_{r-1}$ (since u_r is the supremum over all x). Set

$$y = eT(t_r(x))x \quad \text{so that} \quad T(t)y = eT(t + t_r(x))x \quad \text{for all } t \geq 0$$

By definition of $t_{r+1}(x)$ we have that

$$\|T(t + t_r(x))x\| \leq e^{-r-1} \quad \text{if and only if} \quad t + t_r(x) \geq t_{r+1}(x)$$

Then

$$\|T(t)y\| = e\|T(t + t_r(x))x\| \leq e^{-r} \quad \text{if and only if} \quad t \geq t_{r+1}(x) - t_r(x) = u_r(x)$$

which means that $t_r(y) = u_r(x)$. Now $t_{r-1}(y) = 0$ since $\|T(t' + t_r(x))x\| \leq e^{-r}$ by definition of $t_r(x)$ and so

$$\|T(t')y\| = e\|T(t' + t_r(x))x\| \leq ee^{-r} = e^{-(r-1)} \quad \text{for all } t' \geq 0$$

Hence $u_{r-1}(y) = u_r(x)$. But this means that

$$u_r(x) > u_{r-1} \geq u_{r-1}(y) = u_r(x)$$

is a contradiction. Therefore $u_r \leq u_{r-1}$ for all r as claimed. $\square$

Lemma 5. If the $T(t)$ are (not necessarily proper) contractions (i.e., $\|T(t)\| \leq 1$) then

$$t_r(x) = \sup\{t \geq 0 : \|T(t)x\| \geq e^{-r}\}$$

which is to say the t_r are “stopping times”.

Proof. The $T(t)$ being contractions forces for any $q > 0$ that

$$\|T(t + q)x\| = \|T(q)T(t)x\| \leq \|T(t)x\|\|T(q)\| \leq \|T(t)x\|$$

by the semigroup property. $\square$

5. THE ENTRY TIME GROWTH CHARACTERISTIC $\omega_0^{(ET)}$

Definition 6. The entry time growth characteristic of a C_0 -semigroup $\{T(t)\}$ is defined by

$$\omega_0^{(ET)} := -\left(\lim_r u_r\right)^{-1}$$

which always exists by Lemma 4.

We now show that the entry time growth characteristic is equal to the usual growth characteristic ω_0 defined previously for stable semigroups. Note that if the semigroup is not stable then $\omega_0^{(ET)} = 0$.

Theorem 7. For a stable C_0 semigroup $T(t)$,

$$\omega_0^{(ET)} = \omega_0 = \lim_{t \downarrow 0} \frac{\log \|T(t)\|}{t} = \inf_{t \geq 0} \frac{\log \|T(t)\|}{t} = \lim_{t \rightarrow \infty} \frac{\log \|T(t)\|}{t}$$

Proof. That ω_0 (the usual growth characteristic), which is defined as the limit down to zero above, is equal to the infimum is well-known.

Define $\omega(t) = \frac{1}{t} \log \|T(t)\|$. Then

$$(t+s)\omega(t+s) = \log \|T(t+s)\| \leq \log \|T(t)\| + \log \|T(s)\| = t\omega(t) + s\omega(s)$$

is a subadditive sequence and therefore

$$\lim_{t \rightarrow \infty} \frac{t\omega(t)}{t} = \lim_{t \rightarrow \infty} \omega(t)$$

exists. Call this limit ω . Set $w_r = \omega(t_r)$. By Lemma 3 we know that $w_r = -\frac{r}{t_r}$. Now $\lim_r w_r = \omega$ and

$$-\frac{r+1}{t_{r+1}} = -\frac{r}{t_r} \frac{t_r}{t_{r+1}} - \frac{1}{t_{r+1}}$$

and so

$$w_{r+1} = w_r \frac{t_r}{t_{r+1}} - \frac{1}{t_{r+1}}$$

which means that

$$w_{r+1}t_{r+1} - w_rt_r = -1$$

Consider the case when $\omega > -\infty$. Taking limits in the above, $\lim_r \omega t_r = -1$ and so $\omega_0^{(ET)} = \omega$.

Now consider the case when $\omega = -\infty$. Suppose that $\omega_0^{(ET)} > -\infty$. Then $\lim_r u_r = c > 0$ and so

$$-1 = t_{r+1}w_{r+1} - t_rw_r = t_r(w_{r+1} - w_r) + u_rw_{r+1} \leq u_rw_{r+1}$$

since $w_{r+1} - w_r \leq 0$. But then

$$-1 \leq \lim_r u_rw_{r+1} = c(-\infty) = -\infty$$

contradicting that $\omega_0^{(ET)} > -\infty$. Therefore in either case $\omega_0^{(ET)} = \omega$.

Now we show that $\omega = \omega_0$. Since we know $\omega_0 = \inf_{t \geq 0} \omega(t)$ and $\omega = \lim_{t \rightarrow \infty} \omega(t)$ we already have that $\omega_0 \leq \omega$. For any positive integer n and any $s \geq 0$ we have

$$\omega(ns) = \frac{1}{ns} \log \|T(s)^n\| \leq \frac{1}{ns} \log \|T(s)\|^n = \omega(s)$$

and therefore

$$\lim_{t \rightarrow \infty} \omega(t) = \lim_{n \rightarrow \infty} \omega(ns) \leq \omega(s)$$

which means that

$$\omega = \lim_{t \rightarrow \infty} \omega(t) \leq \inf_{t \geq 0} \omega(t) = \omega_0$$

Therefore $\omega = \omega_0$ and the proof is complete. $\square$

6. EQUIVALENCE OF STABILITY NOTIONS

We now present the main theorems characterizing the various notions of stability. One of our aims in this paper is to collect and clarify the various characterizations of these notions. To this end we include several known characterizations and provide new proofs using our techniques. Specifically, the equivalences in this section, excepting the conditions which involve the relative entry times u_r , are known (see e.g. [Bal05]).

Theorem 8. *For a C_0 -semigroup $\{T(t)\}$ and $\nu > 0$, the following are equivalent:*

- (i) $\{T(t)\}$ is **stable** with **stability index** ν : $(\forall \rho < \nu)(\exists M > 0)(\forall t \geq 0)\|T(t)\| \leq Me^{-\rho t}$
- (ii) $\lim_r u_r = \nu^{-1} < \infty$
- (iii) $\text{spec}(T(t)) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq e^{-t\nu}\}$
- (iv) $\omega_0 = -\nu$

Theorem 9. *If $\{T(t)\}$ is a stable C_0 -semigroup with stability index ν and A is the generator of $T(t)$ with domain D then $\text{spec}(A) \subseteq \{\lambda \in \mathbb{C} : \text{Re } \lambda < -\nu\}$. The converse does not hold.*

Remark. *The spectrum of the generator, $\text{spec}(A)$, depends very delicately on the domain of definition (see the counterexamples below). In particular, the operator A treated as having full domain may well have spectrum much larger than $\text{spec}(A)$.*

Theorem 10. *For a C_0 -semigroup $\{T(t)\}$, the following are equivalent:*

- (i) $\{T(t)\}$ is **superstable** : $\{T(t)\}$ is stable with stability index ∞
- (ii) $\lim_r u_r = 0$
- (iii) $(\forall \nu > 0)(\exists M_\nu > 0)(\forall t \geq 0)\|T(t)\| \leq M_\nu e^{-\nu t}$
- (iv) $T(t)$ are all quasinilpotent: $\text{spec}(T(t)) = \{0\}$
- (v) $\omega_0 = -\infty$

Remark. *The constants M_ν in condition (iii) must tend to infinity as $\nu \rightarrow \infty$ since otherwise the semigroup will be identically 0.*

Theorem 11. *If $\{T(t)\}$ is a superstable C_0 -semigroup with generator A and domain D then $\text{spec}(A) = \emptyset$. The converse does not hold.*

Theorem 12. *For a C_0 -semigroup $\{T(t)\}$ on a Banach space X with generator A having domain D , and $0 \leq k < \infty$ the following are equivalent:*

- (i) $\{T(t)\}$ has **finite time extinction at time** k :
 $(\forall x \in X)(\exists t_\infty(x) \geq 0)(\forall t \geq t_\infty(x))T(t)x = 0$ and $\sup\{t_\infty(x) : \|x\| \leq 1\} = k$
- (ii) $\sum_r u_r = k < \infty$
- (iii) $(\forall \nu > 0)(\exists M_\nu > 0)(\forall t \geq 0)\|T(t)\| \leq M_\nu e^{-\nu t}$ and $\sup_{\nu > 0} \frac{\log M_\nu}{\nu} = k$
- (iv) $(\exists M > 0)(\forall \nu \geq 0)(\forall t \geq 0)\|T(t)\| \leq Me^{-\nu(t-k)}$
- (v) $T(t)$ is nilpotent with nilpotency index k : $T(q) = 0$ for $q > k$ and $T(q) \neq 0$ for $q < k$
- (vi) the resolvent function $R(\lambda, A)$ is entire and $|R(\lambda, A)| \leq C(1 + |\lambda|)^{-N} e^{k|\text{Re } \lambda|}$ for some constants C, N

Remark. *In condition (i), the definition of finite time extinction, the t_∞ can be chosen uniformly on bounded sets, in particular on balls around the origin of finite radius, however, t_∞ cannot be chosen uniformly over x unless the underlying space of the Banach space is compact (i.e., $L^2[0, 1]$ not $L^2[0, \infty)$).*

Theorem 13. *Extinction in finite time implies superstability and superstability implies stability. The converses of both statements are false.*

Theorem 14. *Let $\{T_t\}$ be a C_0 -semigroup. Then, if for some $a > 0$,*

- (i) $\int_a^\infty \|T(t)\|^p dt < \infty$ for some $0 < p < \infty$ *then $\{T(t)\}$ is stable*
- (ii) $\int_a^\infty |\log \|T(t)\||^{-p} dt < \infty$ for some $1 < p < \infty$ *then $\{T(t)\}$ is stable*
- (iii) $\int_a^\infty |\log \|T(t)\||^{-1} dt < \infty$ *then $\{T(t)\}$ is superstable*
- (iv) $\lim_{p \downarrow 0} \int_a^\infty |\log \|T(t)\||^{-p} dt < \infty$ *then $\{T(t)\}$ has finite time extinction*

Remark. *Stability can only occur when $\|T(t)\|$ is eventually bounded by 1 (that is, $t_0 < \infty$). Taking $a = t_0$ will cause the integrals to converge whenever there is some value of a that causes convergence.*

7. PROOFS OF EQUIVALENCES

As usual, let $\{T(t)\}$ be a C_0 -semigroup with generator A having dense domain D on the Banach space X .

Proof. (of Theorem 8). Assume condition (ii) holds:

$$\lim_r u_r = \nu^{-1} < \infty.$$

Fix $\epsilon > 0$. Then there exists r^* such that

$$u_r < \nu^{-1} + \epsilon \quad \text{for all } r \geq r^*.$$

Hence for $r \geq r^*$,

$$t_r - t_{r^*} \leq (r - r^*)(\nu^{-1} + \epsilon).$$

For any $t \geq r^*(\nu^{-1} + \epsilon)$ pick $r \geq r^*$ such that

$$r(\nu^{-1} + \epsilon) + t_{r^*} \leq t < (r + 1)(\nu^{-1} + \epsilon) + t_{r^*}.$$

Since $t \geq t_r$ (as $r(\nu^{-1} + \epsilon) > 0$) we have that $\|T(t)\| \leq e^{-r}$ and

$$t < (r + 1)(\nu^{-1} + \epsilon) + t_{r^*} \quad \text{implies} \quad r > \frac{t - t_{r^*}}{\nu^{-1} + \epsilon} - 1.$$

Then

$$\|T(t)\| \leq e^{-r} < e^{1 - \frac{t - t_{r^*}}{\nu^{-1} + \epsilon}} = e^{1 + \frac{t_{r^*}}{\nu^{-1} + \epsilon}} e^{-t \frac{1}{\nu^{-1} + \epsilon}}$$

So $\{T(t)\}$ has stability index less than $\frac{1}{\nu^{-1} + \epsilon}$. Since ϵ was arbitrary, condition (i) holds.

Conversely, assume (i) holds. Then for any $\rho < \nu$ there is M such that $\|T(t)\| \leq M e^{-\rho t}$. For $t \geq \frac{r - \log M}{\rho}$ we then have that $\|T(t)\| \leq e^{-r}$. Hence

$$t_r \leq \frac{r - \log M}{\rho}$$

Suppose $\lim u_r > \frac{1}{\rho} + 2\delta$ for some $\delta > 0$. Then for sufficiently large r' we have $u_r > \frac{1}{\rho} + \delta$ for $r \geq r'$ so

$$t_{r+1} = t_{r'} + u_{r'} + \cdots + u_r > t_{r'} + (r - r')\frac{1}{\rho} + (r - r')\delta$$

and therefore

$$\frac{r + 1 - \log M}{\rho} - (r - r')\frac{1}{\rho} > (r - r')\delta$$

so

$$\frac{r' + 1 - \log M}{\rho} + r'\delta > r\delta$$

but the lefthand side is constant and the right hand tends to ∞ as $r \rightarrow \infty$. This means that $\lim u_r \leq \frac{1}{\rho}$. Since $\rho < \nu$ is arbitrary we have (ii).

Now assume (iv) holds. Then by Gelfand's spectral radius formula,

$$\sup |spec(T(t))| = \lim_t \|T(t)\|^{\frac{1}{t}} = e^{\omega_0 t} = e^{-t\nu}.$$

Hence (iii) holds. Likewise, if (iii) holds then by Gelfand's formula, $\omega_0 = -\nu$ so (iv) holds.

The equivalence of (i) and (ii) with (iv) is a direct consequence of Theorem 7. This completes the proof that (i) through (iv) are equivalent. $\square$

Proof. (of Theorem 9). This is well-known. $\square$

Proof. (of Theorem 10). The equivalences follow from identical arguments to those for the case of stable semigroups (simply replace ν^{-1} by 0). $\square$

Proof. (of Theorem 11). This follows from Theorem 9: if $z \in \text{spec}(A)$ then the stability index ν is at least $-\text{Re}(z)$ but a superstable semigroup has stability index ∞ . $\square$

Proof. (of Theorem 12). Assume that (ii) holds. Then for any x with $\|x\| \leq 1$ we have that

$$t_{r+1}(x) - t_0(x) = \sum_{j=0}^r u_r(x) \leq \sum_{j=0}^r u_r \rightarrow k$$

hence (i) holds.

Conversely, assume (i) holds and suppose (ii) fails. If $\sum_r u_r = \ell < k$ then

$$t_\infty(x) \leq \ell < k \quad \text{for all } x$$

contradicting (i). So it must be that $\sum_r u_r = \ell > k$. By Lemma 2 we have

$$\sum_r u_r = \sum_r t_{r+1} - t_r = t_\infty - t_0$$

and there is then a sequence x_n such that $t_\infty(x_n) \rightarrow \ell > k$ contradicting (i).

Assume (ii) holds. For $\nu > 0$ pick r^* such that

$$\sup_{r \geq r^*} u_r < \nu^{-1} \quad \text{and set} \quad M_\nu = e^{1+\nu t_{r^*}}.$$

Then, as in the proof of stability, we have

$$r\nu^{-1} \leq t - t_{r^*} < (r+1)\nu^{-1} \quad \text{implies} \quad \|T(t)\| \leq M_\nu e^{-t\nu}.$$

Now

$$\frac{\log M_\nu}{\nu} = \frac{1}{\nu} + t_{r^*}(\nu) \quad \text{and} \quad r^*(\nu) \rightarrow \infty \quad \text{as } \nu \rightarrow \infty$$

hence $\frac{\log M_\nu}{\nu} \rightarrow \sum_r u_r = k$. So (iii) holds.

Assume (iii) holds. Then

$$M_\nu \leq e^{k\nu} \quad \text{for all } \nu$$

hence (iv) holds with $M = 1$.

Assume (iv) holds. Then for $t > k$ we have

$$\|T(t)\| \leq M e^{-\nu(t-k)} \quad \text{for all } \nu \text{ and } t - k > 0$$

so $\lim_\nu e^{-\nu(t-k)} = 0$ hence $\|T(t)\| = 0$. Thus (i) holds.

The equivalence of (i) and (v) is trivial.

To see that (i) and (vi) are equivalent, note that (i) and (vi) both imply $\text{spec}(A) = \emptyset$. Then

$$R(\lambda, A) = \int_0^\infty e^{-\lambda t} T(t) dt \quad \text{for all } \lambda$$

(in general for $\text{Re } \lambda > \omega_0$, see, e.g., [Bal81]).

Hence $R(i\lambda, A)$ is the Fourier Transform of $T(t)$. By the Paley-Wiener Theorem, $R(i\lambda, A)$ is the Fourier Transform of a compactly supported function (i.e. $T(t) = 0$ for all $t \geq k$) if and only if (vi) holds. This argument first appeared in [GK70]. $\square$

8. PROOF OF THE PAZY-TYPE CRITERIA

Lemma 15. *Suppose that $\|T(t)\| \leq 1$ for all $t \geq 0$ (that is, $t_0 = 0$). Let $F : \mathbb{R} \cup \{\pm\infty\} \rightarrow [0, \infty]$ be a decreasing function such that $F(\infty) = 0$. Then*

$$\sum_{r=0}^{\infty} u_r F(r+1) \leq \int_0^\infty F(-\log \|T(t)\|) dt \leq \sum_{r=0}^{\infty} u_r F(r)$$

Proof. Observe that

$$\int_0^\infty F(-\log \|T(t)\|)dt = \int_0^{t_0} F(-\log \|T(t)\|)dt + \sum_{r=0}^\infty \int_{t_r}^{t_{r+1}} F(-\log \|T(t)\|)dt + \int_{t_\infty}^\infty F(-\log \|T(t)\|)dt$$

Now $t_0 = 0$ so the first term on the right is 0. For $t \geq t_\infty$ we have that $\|T(t)\| = 0$ so $F(-\log \|T(t)\|) = F(\infty) = 0$ meaning that the third term on the right is zero.

For the middle terms, note that for $t < t'$,

$$\|T(t')\| = \|T(t)T(t' - t)\| \leq \|T(t)\|\|T(t' - t)\| \leq \|T(t)\|$$

since $\|T(t' - t)\| \leq 1$ by assumption. So

$$\|T(t_{r+1})\| \leq \|T(t)\| \leq \|T(t_r)\| \quad \text{for } t_r \leq t \leq t_{r+1}$$

and therefore, by Lemma 3,

$$-r - 1 = \log \|T(t_{r+1})\| \leq \log \|T(t)\| \leq \log \|T(t_r)\| = -r \quad \text{for } t_r \leq t \leq t_{r+1}$$

and so since F is decreasing

$$F(r + 1) \leq F(-\log \|T(t)\|) \leq F(r) \quad \text{for } t_r \leq t \leq t_{r+1}$$

So for each term in the sum,

$$u_r F(r + 1) = \int_{t_r}^{t_{r+1}} F(r + 1)dt \leq \int_{t_r}^{t_{r+1}} F(-\log \|T(t)\|)dt \leq u_r F(r)$$

□

Proof. (of Theorem 14). First note that if $\limsup \|T(t)\| > 1$ then $u_r = \infty$ for all r so there can be no stability. In this case, none of the three conditions involving integrals can hold. So it is enough to consider the case when $\|T(t)\|$ is eventually bounded by 1. Since

$$\int_0^{t_0} H(\|T(t)\|)dt < \infty$$

for any bounded function H (recall that $t_0 < \infty$ since we have eliminated the other case), we may assume that $\|T(t)\| \leq 1$ for all t : the integral conditions are unaffected by finite translations in time as is the stability of the semigroup.

Recall that condition (i) is the Datko-Pazy Theorem ([Dat70], [Paz72], [Paz83]). Consider the function $F(x) = e^{-px}$ for some fixed $0 < p < \infty$. Then F is decreasing and $F(\infty) = 0$.

By Lemma 15,

$$\sum_{r=0}^\infty u_r e^{-p(r+1)} \leq \int_0^\infty F(-\log \|T(t)\|)dt = \int_0^\infty \|T(t)\|^p dt < \infty$$

For any given r^* observe that since the u_r are nonincreasing,

$$\sum_{r=0}^{r^*-1} u_r e^{-p(r+1)} \geq u_{r^*} \sum_{r=1}^{r^*} e^{-pr}$$

and since $\sum_r e^{-p(r+1)} = C < \infty$,

$$\lim_{r^*} \sum_{r=0}^{r^*-1} u_r e^{-p(r+1)} \geq \lim_{r^*} u_{r^*} C$$

and therefore

$$\lim_r u_r \leq C^{-1} \sum_{r=0}^\infty u_r e^{-p(r+1)} < \infty$$

hence the semigroup is stable.

Condition (ii) is a weakening of the Pazy condition: set $F(x) = x^{-p}$ for the appropriate $1 < p < \infty$. By Lemma 15,

$$\sum_{r=0}^\infty u_r (r+1)^{-p} \leq \int_0^\infty (-\log \|T(t)\|)^{-p} dt < \infty$$

Then, as above, since $\sum_r (r+1)^{-p} = C < \infty$,

$$\lim_r u_r \leq C^{-1} \sum_{r=0}^\infty u_r (r+1)^{-p} < \infty$$

so the semigroup is stable.

Now condition (iii): set $F(x) = \frac{1}{x}$. By Lemma 15,

$$\sum_{r=0}^{\infty} u_r \frac{1}{r+1} \leq \int_0^{\infty} \frac{dt}{-\log \|T(t)\|} < \infty$$

Proceeding as above,

$$\sum_{r=0}^{r^*-1} u_r \frac{1}{r+1} \geq u_{r^*} \sum_{r=1}^{r^*} \frac{1}{r}$$

and since $\sum_{r=1}^{\infty} \frac{1}{r} = \infty$ this means that $\lim_r u_r = 0$ (and in fact converges to 0 faster than the inverse of the harmonic sum $(1 + \dots + \frac{1}{r})^{-1}$). The semigroup is therefore superstable.

Finally condition (iv): set $F_p(x) = x^{-p}$ for $0 < p$. Then, by Lemma 15,

$$\sup_{p>0} \sum_{r=0}^{\infty} u_r (r+1)^{-p} \leq \sup_{p>0} \int_0^{\infty} (-\log \|T(t)\|)^p dt < \infty$$

here we use that $F_p \leq F_{p'}$ for $p \geq p'$ so the hypothesis for $p \downarrow 0$ in fact implies boundedness for all $p > 0$.

Suppose that $\sum_r u_r = \infty$. Then for any K there exists r_K such that $\sum_{r=0}^{r_K-1} u_r \geq K$. Then

$$\sum_{r=0}^{\infty} u_r (r+1)^{-p} \geq \sum_{r=0}^{r_K-1} u_r r_K^{-p} \geq K r_K^{-p}$$

for any $p > 0$ and so

$$\sup_{p>0} \sum_{r=0}^{\infty} u_r (r+1)^{-p} \geq K$$

But K is arbitrary so

$$\infty > \sup_{p>0} \sum_{r=0}^{\infty} u_r (r+1)^{-p} = \infty$$

is a contradiction. The semigroup therefore has finite time extinction. In fact the semigroup goes extinct at time

$$k = \sum_{r=0}^{\infty} u_r = \sup_{p>0} \int_0^{\infty} (-\log \|T(t)\|)^{-p} dt$$

(details here are left to the reader). $\square$

9. COUNTEREXAMPLES

We construct examples of semigroups demonstrating that finite time extinction is strictly stronger than superstability and that superstability is strictly stronger than stability. In particular, we answer a question of Balakrishnan [Bal05] on the existence of superstable semigroups not vanishing in finite time with generator being a differential operator (what he terms a “physical system”). We also remark on a (previously known) example showing that the spectrum of the generator does not fully determine superstability.

9.1. Superstable Without Finite Time Extinction. Consider the Gaussian (probability) measure μ on $\mathbb{R}^+ = [0, \infty)$ given by $d\mu(x) = \sqrt{\frac{2}{\pi}} \exp(-\frac{x^2}{2}) dx$. Let $X = L^2(\mathbb{R}^+, \mu)$. Define the semigroup

$$T(t)f(s) = f(s-t) \text{ for } s \geq t \quad \text{and} \quad T(t)f(s) = 0 \text{ otherwise}$$

on X . The reader may verify that this in fact a semigroup with generator $A = -\frac{d}{ds}$ and domain the appropriate Sobolev space.

Now for $f \in L^2(\mathbb{R}^+, \mu)$ with $\|f\| = 1$ we have

$$\|T(t)f\|^2 = \int_0^{\infty} |f(s-t)|^2 d\mu(s) = \int_0^{\infty} |f(v)|^2 \sqrt{\frac{2}{\pi}} e^{-\frac{(t+v)^2}{2}} dv \leq e^{-\frac{t^2}{2}} \int_0^{\infty} |f(v)|^2 d\mu(v) = e^{-\frac{t^2}{2}}$$

since $e^{-(t+v)^2} \leq e^{-t^2}$ for $t, v \geq 0$. So $\|T(t)\| \leq \exp(-\frac{t^2}{2}) \rightarrow 0$ meaning that the semigroup is superstable. However, $T(t) \neq 0$ for any t .

Taking f to be a norm one (with respect to μ) function concentrated near 0 we see that $\|T(t)\| = \exp(-\frac{t^2}{4})$ and so $t_r = 2\sqrt{r}$ and $u_r = 2(\sqrt{r+1} - \sqrt{r}) \rightarrow 0$ but $\sum u_r = \infty$.

Hence superstability can occur without finite time extinction (even when the generator is merely a derivative). The space $(\mathbb{R}^+, \mu)$ is a variant of the classical Gaussian measure space which arises naturally in the context of stochastic systems and quantum systems, among other areas. Our example can easily be extended to any system with a Gaussian measure (details are left to the interested reader).

9.2. Finite Time Extinction. Define the semigroup

$$T(t)f(s) = f(s+t) \text{ for } s+t \leq 1 \quad \text{and} \quad T(t)f(s) = 0 \text{ otherwise}$$

on $X = \{f \in L^2[0, 1] : f(0) = f(1) = f'(0) = f'(1) = 0\}$. The reader may verify that this is a semigroup with generator $A = \frac{d}{ds}$ and domain Sobolev space. It is clear that $T(t) = 0$ for all $t \geq 1$ so this semigroup has finite time extinction. In fact, $t_r = 1$ for all $r > 0$ so $u_r = 0$ for $r > 0$ and $\sum u_r = 1 < \infty$.

9.3. Stable but Not Superstable. For completeness, we mention the fairly trivial example $T(t)f(s) = e^{-\nu t}f(s)$ (so the generator is $A = \nu I$ and the domain is $\mathcal{D} = L^2$) is clearly stable with index ν . Here t_r is defined by $e^{-\nu t_r} = e^{-r}$ so $t_r = r\nu^{-1}$ meaning $u_r = -\nu^{-1}$.

9.4. Empty Spectrum (for the Generator) but not Superstable. For completeness, we mention an example due to Hille and Phillips [HP57] (chapter 23, section 16).

We first present a superstable semigroup which will be used to develop the actual example of interest. Define $T(t)f(s) := \frac{1}{\Gamma(t)} \int_0^s (s-u)^{t-1} f(u) du$. That this is a semigroup follows from Euler integral identities. The generator A is the derivative of the convolution with log minus a constant: $Af(s) = \frac{d}{ds} \int_0^s \log(s-u) f(u) du - \gamma f(s)$ (γ is Euler's constant). Then $\text{spec}(A) = \emptyset$ and $\|T(t)\| \approx \frac{1}{\Gamma(t)}$. So $\omega_0 = -\infty$ and $T(t) \rightarrow 0$ but $T(t) \neq 0$ for any t . This semigroup is in fact superstable but does not have finite time extinction.

With this construction in hand, we construct the desired example: for $\xi \in \mathbb{C}$ with $\text{Re } \xi > 0$, define $J^\xi f(s) = \frac{1}{\Gamma(\xi)} \int_0^s (s-u)^{\xi-1} f(u) du$. When ξ is taken to be a positive real this yields the semigroup above. There is an analytic extension of J^ξ to ξ purely imaginary. Let $T(t) = J^{it}$. The generator of this semigroup is iA where A is the generator from above. Then $\text{spec}(iA) = \emptyset$ but $0 \notin \text{spec}(T(t))$, i.e. the operators are not quasinilpotent hence not superstable.

The reader is referred to [HP57] for details on these semigroups.

REFERENCES

- [Bal81] A.V. Balakrishnan, *Applied functional analysis*, Springer-Verlag, New York, NY, 1981.
- [Bal05] ———, *Superstability of systems*, Applied Mathematics and Computation **164** (2005), 321–326.
- [Dat70] R. Datko, *Extending a theorem of A. M. Liapunov to Hilbert space*, J. Math. Anal. Appl. **32** (1970), 610–616.
- [GK70] I.C. Gohberg and M.G. Krein, *Theory and applications of Volterra operators in Hilbert space*, American Mathematical Society, Providence, Rhode Island, 1970.
- [HP57] E. Hille and R. Phillips, *Functional analysis and semi-groups*, American Mathematical Society, Providence, Rhode Island, 1957.
- [Lum01] G. Lumer, *On the growth orders of the resolvents for an explicit class of superstable semigroups*, Ulmer Seminare über Funktionalanalysis und Differentialgleichungen **6** (2001).
- [NR92] R. Nagel and F. Răbiger, *Superstable operators on Banach spaces*, Israel J. Math. **81** (1992), no. 1-2, 213–226.
- [Paz72] A. Pazy, *On the applicability of Lyapunov's theorem in Hilbert space*, SIAM J. Math. Anal. **3** (1972), 291–294.
- [Paz83] ———, *Semigroups of operators and applications to partial differential equations*, Springer-Verlag, Berlin, 1983.
- [RW95] F. Răbiger and M. Wolff, *Superstable semigroups of operators*, Indagationes Mathematicae **6** (1995), 481–494.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA: LOS ANGELES, USA

E-mail address: dcreutz@math.ucla.edu

DEPARTMENT OF ELECTRICAL ENGINEERING, UNIVERSITY OF CALIFORNIA: LOS ANGELES, USA

E-mail address: mmazo@ee.ucla.edu

E-mail address: ciprian.preda@feaa.uvt.ro